

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 7 : C11D 3/22, 11/00		A1	(11) International Publication Number: WO 00/42143 (43) International Publication Date: 20 July 2000 (20.07.00)
(21) International Application Number: PCT/EP99/09054 (22) International Filing Date: 24 November 1999 (24.11.99) (30) Priority Data: 9900957.3 15 January 1999 (15.01.99) GB (71) Applicant (for AE AU BB CA CY GB GD GH GM IE IL KE LC LK LS MN MW NZ SD SG SL SZ TT TZ UG ZA ZW only): UNILEVER PLC [GB/GB]; Unilever House, Blackfriars, London EC4P 4BQ (GB). (71) Applicant (for all designated States except AE AU BB CA CY GB GD GH GM IE IL IN KE LC LK LS MN MW NZ SD SG SL SZ TT TZ UG US ZA ZW): UNILEVER NV [NL/NL]; Weena 455, NL-3013 Al Rotterdam (NL). (71) Applicant (for IN only): HINDUSTAN LEVER LIMITED [IN/IN]; Hindustan Lever House, 165/166 Backbay Reclamation, Maharashtra, Mumbai 400 020 (IN). (72) Inventors; and (75) Inventors/Applicants (for US only): SANKHOLKAR, Devadatta, Shivaji [IN/IN]; Hindustan Lever Ltd, Research Centre, Chakala, Andheri (East), Mumbai 400 099 (IN). SIDHESWARAN, Pullimudaliar [IN/IN]; Hindustan Lever Ltd,		(74) Agent: DEKKER, Enno, E., J.; Unilever N.V., Patent Dept., P.O. Box 137, NL-3130 AC Vlaardingen (NL). (81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>	
(54) Title: DETERGENT COMPOSITION			
(57) Abstract A detergent composition suitable for cleaning laundry or hard surfaces comprising up to 50 wt.% of the total detergent composition of detergent active and gum ghatti, a gum of the water swellable, branched hydrocolloids obtained from the species belonging to the genera <i>Anogeissus</i> .			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

- 1 -

Detergent Composition

Technical field

5 The invention relates to a detergent composition for washing fabric or hard surfaces. More particularly the invention relates to compositions for cleaning hard surfaces and for manual dishwashing.

10

Background of the Invention

It is well known that foam volume and foam stability are often used by the consumer to gauge the cleaning efficiency.

15 This is particularly the case when manual cleaning of fabrics and dishes occurs. It is therefore commercially desirable to formulate a detergent composition with improved foaming performance.

20 Commercial hard surface cleaning compositions typically comprise one or more surfactants and a plurality of abrasives dispersed in. Combinations of these together with electrolytes are generally used to form a structuring system as is well known in the art.

25

In the past gums have been incorporated into liquid detergent formulations mainly as viscosity modifiers, as foam stabilisers and to improve product stability.

- 2 -

It has been found that the viscosity of liquid products can be increased by the inclusion of low levels of water-soluble polymers and that the presence of these polymers gives enhanced foaming and detergency.

5

The term Gum as technically employed in industry, refers to plant or microbial polysaccharides or their derivatives that are dispersible in either cold or hot water to produce viscous mixtures or solutions. In general, gums are
10 hydrophobic or hydrophilic high molecular weight molecules, usually with colloidal properties.

EP 124367 (Unilever) discloses the use of selected polymers and Xanthum gum to enhance the foam stability and viscosity
15 of liquid detergents based on dialkyl sulphosuccinates.

Xanthum gum is disclosed as a thickening agent in hard surface cleaners in EP 839907 (Procter and Gamble).

20 The present invention provides a detergent composition that has enhanced foam production, better foam stability and improved detergency.

25 **Description of the Invention**

According to the preferred aspect of the invention there is provided a detergent composition suitable for cleaning laundry or hard surfaces comprising up to 50%wt. of the
30 total detergent composition of detergent active component of gum Ghatti, a gum of the water swellable, branched

- 3 -

hydrocolloids obtained from the species belonging to the genera *Anogeissus*.

According to another aspect of the invention provides a
5 process for the preparation of the detergent composition as claimed in any of the preceding claims comprising the steps of neutralising the active detergent where possible and addition of gum ghatti as an aqueous dispersion to the active detergent.

10

Detailed Description of the Invention

The compositions of the present invention contain as essential components gum ghatti and a detergent. These
15 compositions provide improved cleaning of dishes and laundry in the presence of water, i.e. during use and enhance the foam volume and its stability.

These compositions are of particularly useful for cleaning
20 hard surfaces and in particular for manual dishwashing.

The term dishes as used herein means any utensils involved in food preparation or consumption which may be required to be washed to free them from food particles and other food
25 residues, greases, proteins, starches, gums, dyes and burnt organic residues.

30

- 4 -

Gums :

5 The present invention contains as an essential element gum
Ghatti. Gum Ghatti is derived from *Anogeissus latifolia* a
large gregarious tree belonging to the family Combretaceae,
commonly found in the dry deciduous forests of India and Sri
Lanka. The tree yields a gum or a water swellable, branched
10 hydrocolloid commonly known as Indian gum or *gum ghatti*,
which occurs in straw coloured vermiform tears and dries
without cracking.

The preferred level of gum ghatti in the total composition
15 is from 0.05 to 10 wt% of the total composition, preferably
0.1 to 5 wt%, most preferably 0.1 to 3 wt%.

Optionally other water-soluble polymers such as
hydrophilically substituted polysaccharides or gums such as
20 commonly available vegetable gums, gum arabic, karaya gum,
gellan gum, almond gum, tragacanth gum or Xanthan gum may be
incorporated.

If other water-soluble polymers or gums are present it is
25 preferable if the total level of water-soluble polymer or
gum does not exceed 5 wt% of the total composition and is
preferably is equal to or below 3 wt%.

- 5 -

Detergent Actives:

The composition according to the invention comprise a
5 detergent actives chosen from anionic, nonionic, cationic,
zwitterionic detergent actives or mixtures thereof.

It is preferable if the detergent active comprises an
anionic surfactant. Suitable anionic detergent active
10 compounds are water soluble salts of organic sulphuric
reaction products having in the molecular structure an alkyl
radical containing from 8 to 22 carbon atoms, and a radical
chosen from sulphonic acid or sulphur acid ester radicals
and mixtures thereof.

15
Examples of suitable anionic detergents are sodium and
potassium alcohol sulphates, especially those obtained by
sulphating the higher alcohols produced by reducing the
glycerides of tallow or coconut oil; sodium and potassium
20 alkyl benzene sulphonates such as those in which the alkyl
group contains from 9 to 15 carbon atoms; sodium alkyl
glyceryl ether sulphates, especially those ethers of the
higher alcohols derived from tallow and coconut oil; sodium
coconut oil fatty acid monoglyceride sulphates ; sodium and
25 potassium salts of sulphuric acid esters of the reaction
product of one mole of a higher fatty alcohol and from 1 to
6 moles of ethylene oxide ; sodium and potassium salts of
alkyl phenol ethylene oxide ether sulphate with from 1 to 8
units of ethylene oxide molecule and in which the alkyl
30 radicals contain from 4 to 14 carbon atoms; the reaction
product of fatty acids esterified with isethionic acid and

- 6 -

neutralised with sodium hydroxide where, for example, the fatty acids are derived from coconut oil and mixtures thereof.

- 5 The preferred water-soluble synthetic anionic detergent active compounds are the alkali metal (such as sodium and potassium) and alkaline earth metal (such as calcium and magnesium) salts of higher alkyl benzene sulphonates and mixtures with olefin sulphonates and higher alkyl sulphates, and the higher fatty acid monoglyceride sulphates. The most preferred anionic detergent active compounds are higher alkyl aromatic sulphonates such as higher alkyl benzene sulphonates containing from 6 to 20 carbon atoms in the alkyl group in a straight or branched chain, particular examples of which are sodium salts of higher alkyl benzene sulphonates or of higher-alkyl toluene, xylene or phenol sulphonates, alkyl naphthalene sulphonates, ammonium diamyl naphthalene sulphonate, and sodium dinonyl naphthalene sulphonate. Particularly preferred is Sodium Alkyl Benzene Sulphonate (LAS)

- Suitable nonionic detergent active compounds can be broadly described as compounds produced by the condensation of alkylene oxide groups, which are hydrophilic in nature, with an organic hydrophobic compound which may be aliphatic or alkyl aromatic in nature. The length of the hydrophilic or polyoxyalkylene radical which is condensed with any particular hydrophobic group can be readily adjusted to yield a water-soluble compound having the desired degree of balance between hydrophilic and hydrophobic elements.

- 7 -

Particular examples include the condensation product of aliphatic alcohols having from 8 to 22 carbon atoms in either straight or branched chain configuration with ethylene oxide, such as a coconut oil ethylene oxide

5 condensate having from 2 to 15 moles of ethylene oxide per mole of coconut alcohol; condensates of alkylphenols whose alkyl group contains from 6 to 12 carbon atoms with 5 to 25 moles of ethylene oxide per mole of alkylphenol; condensates of the reaction product of ethylenediamine and propylene

10 oxide with ethylene oxide, the condensate containing from 40 to 80% of polyoxyethylene radicals by weight and having a molecular weight of from 5,000 to 11,000; tertiary amine oxides of structure R_3NO , where one group R is an alkyl group of 8 to 18 carbon atoms and the others are each methyl,

15 ethyl or hydroxyethyl groups, for instance dimethyldodecylamine oxide; tertiary phosphine oxides of structure R_3PO , where one group R is an alkyl group of from 10 to 18 carbon atoms, and the others are each alkyl or hydroxyalkyl groups of 1 to 3 carbon atoms, for instance

20 dimethyldodecylphosphine oxide; and dialkyl sulphoxides of structure R_2SO where the group R is an alkyl group of from 10 to 18 carbon atoms and the other is methyl or ethyl, for instance methyltetradecyl sulphoxide; fatty acid alkylolamides; alkylene oxide condensates of fatty acid

25 alkylolamides and alkyl mercaptans.

Suitable amphoteric detergent-active compounds that optionally can be employed are derivatives of aliphatic secondary and tertiary amines containing an alkyl group of 8

30 to 18 carbon atoms and an aliphatic radical substituted by an anionic water-solubilizing group, for instance sodium 3-

- 8 -

dodecylamino-propionate, sodium 3-dodecylaminopropane
sulphonate and sodium N-2-hydroxydodecyl-N-methyltaurate.
Suitable cationic detergent-active compounds are quaternary
ammonium salts having an aliphatic radical of from 8 to 18
5 carbon atoms, for instance cetyltrimethyl ammonium bromide.

Suitable zwitterionic detergent-active compounds that
optionally can be employed are derivatives of aliphatic
quaternary ammonium, sulphonium and phosphonium compounds
10 having an aliphatic radical of from 8 to 18 carbon atoms and
an aliphatic radical substituted by an anionic water-
solubilising group, for instance 3-(N-N-dimethyl-N-
hexadecylammonium) propane-1-sulphonate betaine, 3-
(dodecylmethyl sulphonium) propane-1-sulphonate betaine and
15 3-(cetylmethylphosphonium) ethane sulphonate betaine.

Further examples of suitable detergent-active compounds are
compounds commonly used as surface-active agents given in
the well-known textbooks "Surface Active Agents", Volume I
20 by Schwartz and Perry and "Surface Active Agents and
Detergents", Volume II by Schwartz, Perry and Berch.

The total amount of detergent active compound to be employed
in the detergent composition of the invention will generally
25 be from 1.5 to 25%, preferably from 2 to 15% by weight.

Abrasives:

A particulate abrasive phase is a useful ingredient of
compositions according to the present invention.

30

- 9 -

Preferably, the particulate phase comprises a particulate abrasive that is insoluble in water. In the alternative, the abrasive may be soluble and present in such excess to any water present in the composition that the solubility of the
5 abrasive in the aqueous phase is exceeded and consequently solid abrasive exists in the composition.

Suitable abrasives can be selected from, particulate zeolites, calcites, dolomites, feldspars, silicas,
10 silicates, other carbonates, aluminas, bicarbonates, borates, sulphates and polymeric materials such as polyethylene.

Preferred abrasives for use in general purpose compositions
15 have Mohs hardness 2-6 although higher hardness abrasives can be employed for specialist applications.

Preferred average particle sizes for the abrasive fall in the range 0.5-400 microns, with values of around 10-200
20 microns being preferred.

Preferred levels of abrasive range from 4-95wt % on product, more preferably in the range 20-60wt%. The physical form of the product will be influenced by the level of abrasive
25 present.

The most preferred abrasives are calcium carbonate (as Calcite), mixtures of calcium and magnesium carbonates (as dolomite), sodium hydrogen carbonate, potassium sulphate,
30 zeolite, alumina, hydrated alumina, feldspars, talc and silica.

- 10 -

Calcite, feldspar and dolomite and mixtures thereof are particularly preferred due to their low cost, suitable hardness and colour.

5

Builders:

Compositions of the invention may contain a detergency builder.

10

The detergency builders used in the formulation are preferably inorganic and suitable builders include, for example, alkali metal aluminosilicates (zeolites), sodium carbonate, sodium tripolyphosphate (STPP), tetrasodium pyrophosphate (TSPP), and combinations of these.

15

Typical examples of phosphorus-containing inorganic builders, when present, include the water-soluble salts, especially alkali metal pyrophosphates, orthophosphates and polyphosphates. Specific examples of inorganic phosphate builders include sodium and potassium tripolyphosphates, pyrophosphates and hexametaphosphates.

20

Suitable examples of non-phosphorus-containing inorganic builders, when present, include water-soluble alkali metal carbonates, bicarbonates, sesquicarbonates, borates, silicates, including layered silicates such as SKS-6 ex. Hoechst, metasilicates, and crystalline and amorphous aluminosilicates. Specific examples include sodium carbonate (with or without calcite seeds), potassium carbonate, sodium and potassium bicarbonates, silicates including layered silicates and zeolites. The preferred

30

- 11 -

sodium aluminosilicates of this type are the well-known commercially available zeolites 4A and X, maximum aluminium zeolite P (zeolite MAP) and mixtures thereof.

- 5 It is preferred for hand dish wash formulations if the builder material is, sodium tripolyphosphate (STPP)

Builders are suitably used in an amount ranging from 0.1 to
10 50% by wt, preferably from 0.5 to 30% by wt, most preferably from 0.5 to 10 wt%.

Fillers:

- 15 Fillers suitable for use in the formulation include kaolin, calcium carbonate (calcite), talc, soapstone, china clay and the like, used singly or in combination, suitably in an amount ranging from 10 to 75% by weight, preferably from 30 to 70 wt%.

20

Solvents:

- Solvents suitable in the compositions of the present invention are Alkanolamines, which can be mono- or poly-
25 functional as regards the amine and hydroxy moieties. Preferred alkanolamines are generally of the formulation H_2N-R_1-OH where R_1 is a linear or branched alkyl chain having 2-6 carbons. Particularly preferred alkanolamine especially to clean tough or aged soil is 2-amino-2-methyl-1-propanol
30 (AMP) or monoethanol amine.

- 12 -

Other suitable solvents include saturated and unsaturated, linear or branched hydrocarbons, and/or materials of the general formula:



wherein R_1 and R_2 are independently C1-7 alkyl or H, but not both hydrogen, m and n are independently 0-5. Suitable glycol ethers include di-ethylene glycol mono n-butyl ether, 10 mono-ethylene glycol mono n-butyl ether, propylene glycol n-butyl ether and mixtures thereof. Typical levels of solvent range from 1-15% wt.

Electrolyte base:

15 Suitable electrolyte bases include soluble carbonates and bicarbonates, although use of hydroxides and other alkaline salts is not excluded. Alkali metal carbonates are particularly preferred, with potassium carbonate being the most preferred. Typical levels of electrolytes range from 20 0.5-5%wt, with 1-2.5%wt being particularly preferred. The level of the electrolyte should be such that in use the pH of the composition is raised above the pKa of the alkanolamine, and preferably to a pH at least one unit above 25 the pKa of the alkanolamine.

Rheology and Structuring Agents:

As described above the compositions of the invention be in 30 the form of solid, liquids, pastes or gels. Other suitable rheological control agents can be present especially when

- 13 -

the compositions contain significant amounts of water or low viscosity surfactants.

5 **Other Ingredients:**

Other ingredients such as perfumes, colouring agents, fluoresces, enzymes and bleaches can also be used in the formulation.

10

Product Form:

Detergent compositions of the present may formulated in different forms such as powders, granules, bars, liquids, pastes and gels. However it is preferable if the product is in solid form that is a powder, granule, bar or paste. It is most preferable if the composition is a bar.

20 The invention will now be illustrated with respect to the following non-limiting examples.

Examples:

25 Process for preparation of the composition:

The formulations disclosed in Table 1 were prepared using conventional bar processing technology. The ingredients were mixed in a sigma mixer, extruded into bars and then cut into billets and stamped. In the experimental examples 2 to 5 the gum ghatti dispersed in water was incorporated.

30

Table 1

Composition (%wt)	Ex 1	Ex 2	Ex 3	Ex 4	Ex 5
Na LAS	7	7	7	7	7
China clay	24.1	24.1	24.1	24.1	24.1
Feldspar	29.0	29.0	29.0	29.0	29.0
Aluminium sulphate	2.0	2.0	2.0	2.0	2.0
Alkaline silicate	3.5	3.5	3.5	3.5	3.5
Soda	6.7	6.7	6.7	6.7	6.7
Calcium hydroxide	1.0	1.0	1.0	1.0	1.0
STPP	1.0	1.0	1.0	1.0	1.0
Calcite	11.0	11.0	11.0	11.0	11.0
Gum Ghatti	-	0.25	0.5	2	6.0
Minor ingredients	0.2	0.2	0.2	0.2	0.2
Water	to 100	to 100	to 100	to 100	to 100

5 In use properties of the bar:

a. **Tough soil cleaning:**

The mobile soil that has been polymerised by heating at high temperature is known as tough soil. When the soil is smeared
 10 uniformly on a stainless steel plate and heated it forms a polymerised film on the plate.

- 15 -

0. 2g of the product as defined in the different Examples mentioned in Table 1 were uniformly rubbed on the soiled plate and then rinsed with water. The amount of soil removed was evaluated gravimetrically. The experiment was conducted
5 with replicates and analysed statistically.

b. Lather Measurements:

0.1g of the detergent bars Examples 1 to 5 were dissolved in 20 ml water were taken in different graduated measuring
10 cylinders. The cylinders were shaken 30 times and the volume of lather and the nature of lather were noted.

c. Cleaning of unpolymerised oily soil:

Uniform quantity of vegetable oil is spread on stainless
15 steel plates and the number of such soiled plates that can be cleaned for a particular quantity of the product was assessed using samples according to Examples 1-5.

In all the above evaluations 2% *gum ghatti* dispersed in
20 water was used for comparison, since at the corresponding level of dilution of the gum dispersion as done for the detergent bar, the lather, % cleaning of tough soil and number of plates cleaned are negligible or zero.

25 The data presented in Table 2 shows that the incorporation of gum at different levels into the detergent bar gave a very creamy lather and significant improvement in tough soil cleaning as well as increase in number of plates that could be cleaned. It was also found that the bar processability
30 improved significantly by the incorporation of the gum into

- 16 -

the bar but at high levels (6%) of incorporation, the formulation became pasty and difficult to process.

Table 2

5

Composition	Lather Volume (ml) Characteristics		% Cleaning of tough soil	Number of plates cleaned
Example 1	97	Thin & Unstable	21.4	5
Example 2	103	Slight creamy	26.7	6
Example 3	103	Creamy	29.1	6
Example 4	102	Creamy	30.2	6
Example 5	90	Very thick	28.5	7
2% <i>gum ghatti</i> dispersion	5	Creamy	3.4	0

Comparison Of *gum ghatti* with *gum Arabic*:

- 10 Detergent bars were prepared according to Example 1, Example 3 and in Example 6 the 0.5% of *gum arabic* was incorporated in place of *gum ghatti*. These samples were tested by the procedure described above for lather measurement, tough soil cleaning and the cleaning of unpolymerised soil. The data
- 15 presented in Table 3 shows that *gum ghatti* was significantly superior to *gum Arabic* in lather volume and tough soil cleaning. In addition, the processability of the bar with *gum ghatti* was significantly superior to *gum Arabic*.

20

Table 3

Composition	Lather		% cleaning of tough soil	Number of plates cleaned
	Volume (ml)	Characteristics		
Example 1	97	Thin & Unstable	21.4	5
Example 3	103	Creamy	29.1	6
Example 6	86	Creamy	28.3	6

5

It is thus possible to obtain a synergistic detergent composition by incorporating the gum or the water swellable, branched hydrocolloids obtained from the species belonging

10 to the genera *Anogeissus*.

CLAIMS

1. A detergent composition suitable for cleaning laundry or hard surfaces comprising up to 50%wt. of the total detergent composition of detergent active and gum ghatti, a gum of the water swellable, branched hydrocolloids obtained from the species belonging to the genera *Anogeissus*.
5
2. A detergent composition according to claim 1 in which the level of gum ghatti is from 0.1 to 5 wt% of the total composition.
10
3. A detergent composition according to claim 1 or claim 2 in which the composition further comprises a particulate abrasive.
15
4. A detergent composition according to claim 3 in which the level of particulate abrasive is from 20 to 60 wt% of the total composition.
20
5. A detergent composition according to claim 3 or claim 4 in which the particulate abrasive is selected from calcite, feldspar, dolomite or mixtures thereof.
25
6. A detergent composition according to any preceding claim in which the detergent active is an anionic surfactant.
7. A detergent composition according to any preceding claim in which the detergent active is sodium alkyl benzene sulphonate
30

- 19 -

8. A detergent composition according to any preceding claim in which the total level of detergent active is from 1.5 to 25 wt%.
- 5 9. A detergent composition according to any preceding claim which further comprises a detergency builder.
- 10 10. A process for the preparation of the detergent composition as claimed in any of the preceding claims comprising the steps of neutralising the active detergent where possible and addition of gum ghatti as an aqueous dispersion to the active detergent.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 99/09054

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C11D3/22 C11D11/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C11D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DE 195 26 033 A (HENKEL KGAA) 23 January 1997 (1997-01-23) claims 1-7	1,2,6-8
X A	US 4 581 042 A (WILLMORE GEORGE A) 8 April 1986 (1986-04-08) column 3, line 15 - line 28; claims 1,5,7,10	1,3,5-7, 9,10 2,4,8
X	DATABASE WPI Section Ch, Week 199617 Derwent Publications Ltd., London, GB; Class D25, AN 1996-167424 XP002132873	1,6
A	& JP 08 048999 A (NISSHIN OIL MILLS LTD), 20 February 1996 (1996-02-20) abstract	2
	-/-	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

13 March 2000

Date of mailing of the international search report

23/03/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Saunders, T

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 99/09054

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	US 5 942 478 A (LOPES JOHN A) 24 August 1999 (1999-08-24) claims 1,8,16	1,7-9
A	EP 0 124 367 A (UNILEVER NV) 7 November 1984 (1984-11-07) cited in the application claim 1; example 14	1,6-9

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No
PCT/EP 99/09054

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE 19526033 A	23-01-1997	NONE	
US 4581042 A	08-04-1986	NONE	
JP 8048999 A	20-02-1996	JP 2837360 B	16-12-1998
US 5942478 A	24-08-1999	NONE	
EP 0124367 A	07-11-1984	AT 31939 T	15-01-1988
		AU 552499 B	05-06-1986
		AU 2721084 A	01-11-1984
		BR 8401935 A	04-12-1984
		CA 1224107 A	14-07-1987
		DK 212584 A	30-10-1984
		GB 2140024 A, B	21-11-1984
		GR 81900 A	12-12-1984
		IN 159974 A	13-06-1987
		JP 59207995 A	26-11-1984
		NO 841612 A	30-10-1984
		NZ 207944 A	12-11-1986
		PH 19177 A	23-01-1986
		PT 78501 A, B	01-05-1984
		US 4576744 A	18-03-1986
		ZA 8403161 A	24-12-1985